

new 339

UNIVERSIDADE DE SÃO PAULO
Instituto de Ciências Matemáticas e de Computação
ISSN 0103-2577

**A NOTE ON THE RELATIONSHIP BETWEEN SPECTRAL
RADIUS AND NORMS OF BOUNDED LINEAR OPERATORS**

HILDEBRANDO M. RODRIGUES
J. SOLÀ-MORALES

Nº 292

NOTAS

Série Matemática



São Carlos – SP
Abr./2008

SYSNO	1661079
DATA	1/1
ICMC - SBAB	

Resumo:

Seja X um espaço de Banach e $\mathcal{L}(X)$ a álgebra de Banach dos operadores lineares limitados em X . Nesta nota provamos que se temos um subconjunto compacto K de uma sub-álgebra comutativa de $\mathcal{L}(X)$, dado $\varepsilon > 0$, então é possível definir uma nova norma em X , equivalente a sua dada norma, de tal modo que dentro de uma vizinhança U_ε desse conjunto compacto na subálgebra, as normas de todos os operadores diferem de seus respectivos raios espectrais a menos de ε . Se X for um espaço de Hilbert então é possível definir esta nova norma como um norma Hilbertiana.

A NOTE ON THE RELATIONSHIP BETWEEN SPECTRAL RADIUS AND NORMS OF BOUNDED LINEAR OPERATORS.

HILDEBRANDO M. RODRIGUES[†] AND J. SOLÀ-MORALES[‡]

ABSTRACT. Let X be a Banach space and $\mathcal{L}(X)$ be the Banach algebra of bounded operators on X . In this note we prove that if we have a compact subset K of a commutative sub-algebra of $\mathcal{L}(X)$, and given $\varepsilon > 0$, then it is possible to define a new norm in X , equivalent to its given norm, in such a way that inside a neighborhood U_ε of this compact set in the sub-algebra, the norms of all the operators differ from their spectral radius in less than ε . If X is a Hilbert space then it is possible to define this new norm as an Hilbertian norm.

Mathematics Subject Classification Primary 47A10, 47A30; Secondary 46J05

keywords spectral theory, adapted norms, spectral radius.

When one wants to estimate the norms of functions of bounded operators the properties of norms on $\mathcal{L}(X)$ usually are very helpful. But imposing assumptions on the spectral radius of an operator is better than imposing on its norm, because the first is intrinsic and the second one is not. But for certain sets of bounded operators one can change the norm of the space to an equivalent norm in such a way that the norms of the bounded operators are as near as

Date: April 24, 2008.

[†]Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo-Campus de São Carlos, Caixa Postal 668, 13560-970 São Carlos SP, Brazil

e-mail: hmr@icmc.usp.br.

[‡]Departament de Matemàtica Aplicada 1, Universitat Politècnica de Catalunya, Av. Diagonal 647, 08028 Barcelona, Spain, e-mail: jc.sola-morales@upc.edu.

[†]Partially supported by CNPq Proc.: 301994/85-4, Programa Pronex-Projeto Tematico, CNPq-FAPESP n. 2003/10042-0, FAPESP Proc.: 2005/55694-0, Projeto Milenio/IMPA Proc.: 420108/2005-0 and CAPES Proc.: 1356062, Brasil.

[‡]Partially supported by MEC, Spain (MTM2005-07660-C02-01 and PHPB2006-006) and by DURSI, Generalitat de Catalunya (2005BE00245).

we wish to their spectral radius. This fact has been widely used by many authors, including C. Pugh classical paper [1], for example.

In our previous papers Rodrigues-Sola Morales [2] and [3] we also used this fact, and in [2] we proved that if we have a finite family of bounded linear operators T_n , such that T_n commutes with T_m for every m, n , then it is possible to define a new norm on the Banach space X , equivalent to the initial norm, in such a way that in the new norm the spectral radius, $r(T_n)$, approximates the norm of the operator T_n , for every n . For just one operator this result was proved by Holmes [4]. When the family consists of two operators, and of the form T, T^{-1} , the result had already appeared in the mathematical literature, like for example in Cabre, Fontich, de la Llave [6].

In the present note we improve the above result by proving that if we have a compact subset, K , of a commutative sub-algebra of $\mathcal{L}(X)$, and given $\varepsilon > 0$, then it is possible to define a new norm on X , equivalent to its given norm, in such a way that in a neighborhood U_ε of this compact set inside the sub-algebra, the norms of all the operators differ from their spectral radius in less than ε . If X is a Hilbert space then it is possible to define a new Hilbertian norm with the specified property. This answers a question posed by Charles Pugh to one of the authors.

In a Banach space X with a norm $|\cdot|$, given an operator $T \in \mathcal{L}(X)$ and a number $\varepsilon > 0$ one can define a new norm $|\cdot|_{T,\varepsilon}$ with the formula:

$$|x|_{T,\varepsilon} = \left[|x|^2 + \left(\frac{|Tx|}{M} \right)^2 + \cdots + \left(\frac{|T^m x|}{M^m} \right)^2 \right]^{1/2},$$

where $M = r(T) + \varepsilon$ and $m \in \mathbb{N}$ has been chosen as the first integer such that $|T^m|^{1/m} < M$, where we denote by $r(T)$ the spectral radius of T . It is easy to see that this norm is equivalent to the original norm and that

$$|T|_{T,\varepsilon} < r(T) + \varepsilon.$$

We are interested in pointing out the following property, that plays a key role in the rest of this paper:

Lemma 1. *If $T, S \in \mathcal{L}(X)$ and $ST = TS$, then*

$$|S|_{T,\varepsilon} \leq |S|.$$

Proof: It is easy to see that $|Sx|_{T,\varepsilon} \leq |S||x|_{T,\varepsilon}$, for every $x \in X$.

As a consequence of this simple lemma, one can prove the following:

Proposition 1. *If $T, S \in \mathcal{L}(X)$ and $ST = TS$, then*

- (a) $r(T + S) \leq r(T) + r(S)$ and $r(TS) \leq r(T) r(S)$.
- (b) *The spectral radius of $T \in \mathcal{L}(X)$ changes continuously when T is subjected to a small change which commutes with T .*

Proof: (a) Given $\varepsilon > 0$, from Lemma 1, there exists a new norm $\|\cdot\|$, such that $\|T\| < r(T) + \varepsilon$, $\|S\| < r(S) + \varepsilon$, So

$$r(T + S) \leq \|T + S\| \leq \|T\| + \|S\| < r(T) + r(S) + 2\varepsilon$$

$$r(TS) \leq \|TS\| \leq \|T\|\|S\| < r(T)r(S) + \varepsilon(r(T) + r(S)) + \varepsilon^2.$$

(b) From (a) it follows that:

$$r(T) = r(T - S + S) \leq r(T - S) + r(S) \leq \|T - S\| + r(S)$$

and

$$r(S) = r(S - T + T) \leq r(S - T) + r(T) \leq \|T - S\| + r(T)$$

Therefore $-\|T - S\| \leq r(T) - r(S) \leq \|T - S\|$. This implies the continuity.

The following is our main result:

Theorem 1. *Let X be a Banach space with a norm $|\cdot|$ and let K be a compact subset of $\mathcal{L}(X)$ such that $TS = ST$ for all $T, S \in K$.*

- (i) *Given $\varepsilon > 0$ it is possible to define a new norm $\|\cdot\|$ equivalent to the given norm $|\cdot|$ in such a way that*

$$\|T\| < r(T) + \varepsilon, \tag{0.1}$$

for every $T \in K$.

- (ii) Given a set $\mathcal{U}$, not necessarily compact, with $K \subset \mathcal{U} \subset \mathcal{L}(X)$ and such that $TS = ST$ for all $T, S \in \mathcal{U}$, property (0.1) also holds for T in a neighborhood U_ε of K inside $\mathcal{U}$.

Proof: For all $T \in K$ we consider the open ball centered at T of radius $\varepsilon/3$. This is a covering of K , from which one can extract a finite covering, with centers $T_1, T_2, \dots, T_n$.

By using the above procedure, we modify successively the norm $|\cdot|$ of X to $|\cdot|_{T_1, \varepsilon/3}$, then by using the operator T_2 we modify the norm $|\cdot|_{T_1, \varepsilon/3}$ and so on.

At the end, we have an equivalent norm $\|\cdot\|$, such that, by Lemma 1,

$$\|T_i\| < r(T_i) + \varepsilon/3,$$

for $i = 1, 2, \dots, n$ and such that $\|S\| \leq |S|$ for all $S \in \mathcal{U}$.

Given now any $T \in K$, there exists a T_j such that $|T - T_j| < \varepsilon/3$, and we have:

$$\|T\| \leq \|T_j\| + \|T - T_j\| \leq \|T_j\| + |T - T_j|,$$

where we have used that $T - T_j$ commutes with all of the T_i 's. And now

$$\|T\| \leq \|T_j\| + \varepsilon/3 < r(T_j) + \varepsilon/3 + \varepsilon/3.$$

Finally, by Proposition 1,

$$r(T_j) \leq r(T) + r(T - T_j) \leq r(T) + |T - T_j| < r(T) + \varepsilon/3$$

Then $\|T\| < r(T) + \varepsilon$ and the proof of (i) is complete.

To prove part (ii) we first use part (i) and consider the norm $\|\cdot\|$ such that

$$\|T\| < r(T) + \varepsilon$$

for all $T \in K$. Then we observe that because of Proposition 1 the map $T \mapsto r(T)$ is continuous from $\mathcal{U}$ to $\mathbb{R}$, where $\mathcal{U}$ inherits the topology of the norm $\|\cdot\|$ of X . Then it is obvious that the set

$$\{T \in \mathcal{U} : \|T\| - r(T) < \varepsilon\}$$

contains K and is an open set in the relative topology of $\mathcal{U}$.

Remark 1. The previous results hold true without changing to a new norm if X is a Hilbert space and T is a selfadjoint or a normal operator.

Remark 2. The result (ii) needs not to be true if one drops the property that $TS = ST$ for all $T, S \in \mathcal{U}$, that is if we speak about the neighborhoods of K in X . As it is explained in Kato [5] the spectral radius can shrink discontinuously in arbitrary neighborhoods of an operator.

Remark 3. Note that every compact $K \subset \mathcal{L}(X)$ as in Theorem 1 generates a commutative Banach algebra $\mathcal{A} \subset \mathcal{L}(X)$, so the set $\mathcal{U}$ could be taken as $\mathcal{U} = \mathcal{A}$.

Let us explain some more what could be one of the uses of Theorem 1 above. If X is a Banach space, and $L \in \mathcal{L}(X)$, then one can consider the operator $f(L)$ defined with the Operational Calculus formula:

$$f(L) = \int_{\gamma} f(z)(L - zI)^{-1} dz. \quad (0.2)$$

The typical case is when f is an holomorphic function in an open set $\Omega \subset \mathbb{C}$ with $\sigma(L) \subset \Omega$ and $\gamma : [0, 1] \rightarrow \Omega \setminus \sigma(L)$ is a closed rectifiable curve. Then, the set

$$K = \{(L - z)^{-1} | z \in \{\gamma\}\} \subset \mathcal{L}(X)$$

is a compact set of commuting operators.

Then the previous Theorem can be applied, and given an arbitrary $\varepsilon > 0$ there exists a suitable new equivalent norm $\|\cdot\|$ such that

$$\|f(L)\| \leq \int_0^1 |\gamma'(s)| \frac{|f(\gamma(s))|}{\text{dist}(\gamma(s), \sigma(L))} ds + \varepsilon \text{ length } [\gamma], \quad (0.3)$$

an inequality that can be useful in many cases. And we point out that the inequality (0.3), and inequalities of similar type, can be derived from (0.2) without requiring $f(z)$ to be holomorphic in Ω containing σL . The only thing that one needs is $f(\gamma(s))$ to be integrable in $(0, 1)$.

Some of the above results can be extended to abstract Banach algebras.

Theorem 2. *Let $\mathcal{A}$ be a commutative algebra of Banach with norm $|\cdot|$. Let K be a compact subset of $\mathcal{A}$. Then given $\varepsilon > 0$, there exists a new norm $\|\cdot\|$ in $\mathcal{A}$ equivalent to $|\cdot|$ and a neighborhood U_ε of K , such that for every $T \in U_\varepsilon$*

$$\|T\| < r(T) + \varepsilon.$$

Proof: The proof follows the ideas of Theorem 1 with minor changes. We first observe that $\mathcal{A}$ with norm $|\cdot|$ is a Banach space. For $T \in \mathcal{A}$, we define a new Banach space norm as:

$$|S|_{T,\varepsilon} = \left[|S|^2 + \left(\frac{|ST|}{M} \right)^2 + \cdots + \left(\frac{|S^m T|}{M^m} \right)^2 \right]^{1/2},$$

where M and m are given as above. This norm will induce a Banach algebra norm, $|\cdot|_{T,\varepsilon}$, in $\mathcal{L}(\mathcal{A})$, defined in the usual way and $\mathcal{A} \subset \mathcal{L}(\mathcal{A})$.

It is possible to prove that this norm restricted to $\mathcal{A}$ is equivalent to the norm $|\cdot|$.

The remaining part of the proof follows the steps of the proof of Theorem 1.

REFERENCES

- [1] C. C. Pugh, *On a Theorem of P. Hartman*, American Journal of Mathematics **91**, 2, (1969), 363-367.
- [2] H. M. Rodrigues and J. Solà-Morales, *Linearization of Class C^1 for Contractions on Banach Spaces*, J. Differential Equations **201**, 2, (2004), 351-382.
- [3] H. M. Rodrigues and J. Solà-Morales, *Smooth Linearization for a saddle on Banach Spaces*, J. Dynamics and Differential Equations **16**, 3, (2004), 767-793.
- [4] R. B. Holmes, *A formula for the spectral radius of an operator*, Amer. Math. Monthly **75** (1968), 163-166.
- [5] Kato, T. (1976). *Perturbation Theory for Linear Operators*, Second Edition, Springer Verlag, 1976.
- [6] X. Cabré, E. Fontich and R. de la Llave *The parametrization method for invariant manifolds, I. Manifolds associated to non-resonant subspaces*. Indiana Univ. Math. J. **52**, 2, (2003), 283-328.

NOTAS DO ICMC

SÉRIE MATEMÁTICA

- 291/2008 MATTOS, D.; SANTOS, E. L. - On nonsymmetric theorems for (H, G) -coincidences.
- 290/2008 BAPTISTELLI, P.H.; MANOEL, M. - The σ -isotypic decomposition and the σ -index of reversible-equivariant systems.
- 289/2007 CARVALHO, A.N.; DLOTKO, T.; NASCIMENTO, M.J.D. - Non-autonomous semilinear evolution equations with almost sectorial operators.
- 288/2007 CARVALHO, A.N.; DLOTKO, T. - Dynamics of the viscous Cahn-Hilliard equations.
- 287/2007 BIASI, C.; MATTOS, D.; SANTOS, E.L. - Applications of the non-standard Borsuk-Ulam theorem.
- 286/2007 NABARRO, A.C.; TARI, F. - Families of surfaces and conjugate curve congruences.
- 285/2007 FEDERSON, M. - Converse Lyapunov theorems for retarded functional differential equations.
- 284/2007 ANTONELI, F.; DIAS, A. P.; BAPTISTELLI, P. H.; MANOEL, M. G. - Invariant theory and reversible-equivariant vector fields.
- 283/2007 TAHZIBI, A. - Ergodic properties of dynamical systems beyond uniform hiperbolicity (38th Iranian Mathematical Conference) Zanzan-Iran.
- 282/2007 LLIBRE, J.; OLIVEIRA, R.D.S. - Phase portraits of quadratic polynomial vector fields having a rational first integral of degree 3.