

UNIVERSIDADE DE SÃO PAULO
Instituto de Ciências Matemáticas e de Computação
ISSN 0103-2577

FINITE-DIMENSIONAL GLOBAL ATTRACTORS IN BANACH SPACES

ALEXANDRE N. CARVALHO
JOSÉ A. LANGA
JAMES C. ROBINSON

Nº 306

NOTAS

Série Matemática



São Carlos – SP
Fev./2009

SYSNO	1729078
DATA	1 1
ICMC - SBAB	

Resumo. *Obtemos cotas superiores para a dimensão fractal de subconjuntos negativamente invariantes de espaços de Banach, um problema que pode facilmente ser reduzido ao recobrimento da imagem da bola unitária sob uma transformação linear por uma coleção de bolas de raio menor. Como uma aplicação da teoria abstrata, mostramos que os atratores globais de uma classe bastante ampla de problemas parabólicos semilineares têm dimensão finita.*

FINITE-DIMENSIONAL GLOBAL ATTRACTORS IN BANACH SPACES

ALEXANDRE N. CARVALHO¹, JOSÉ A. LANGA², AND JAMES C. ROBINSON³

ABSTRACT. Negatively invariant sets, Fractal dimension We provide bounds on the upper box-counting dimension of negatively invariant subsets of Banach spaces, a problem that is easily reduced to covering the image of the unit ball under a linear map by a collection of balls of smaller radius. As an application of the abstract theory we show that the global attractors of a very broad class of parabolic partial differential equations (semilinear equations in Banach spaces) are finite-dimensional.

1. INTRODUCTION

There are now many techniques available for bounding the (box-counting) dimension of the global attractors that have been shown to exist for a variety of interesting models of mathematical physics (for many examples see Temam, 1988).

To our knowledge, the earliest such result is due to Mallet-Paret (1976), who showed that if K is a compact subset of a Hilbert space H , $f : H \rightarrow H$ is a continuously differentiable map, $f(K) \supset K$ (' K is negatively invariant'), and the derivative of f is everywhere equal to the sum of a compact map and a contraction, then the upper box-counting dimension of K is finite ($\dim_B(K) < \infty$). This method was subsequently generalised by Mañé (1981) to treat subsets of Banach spaces.

Both these methods rely on approximating the image of any ball B under f as the sum of a ball in a finite-dimensional subspace of X plus a 'small' error. Douady & Oesterlé (1980) developed a method for subsets of $\mathbb{R}^n$ that instead approximates $f(B)$ by an ellipse, plus an error. This approach, extended to subsets of Hilbert spaces by Constantin & Foias (1985), produces significantly improved bounds on the dimension and remains the most powerful to date.

We concentrate here on the case of subsets of Banach spaces, since the Hilbert space theory has been well developed. The problem reduces, as we shall see, to bounding the number of balls required to cover the image of the unit ball under a certain family of linear maps.

We choose to work with the upper box-counting dimension ($\dim_B(\cdot)$) for two main reasons. First, it provides an upper bound on the topological ($\dim_T(\cdot)$) and Hausdorff dimension ($\dim_H(\cdot)$) (see Hurewicz & Wallman (1941) for more on the topological dimension and a proof that $\dim_T(\cdot) \leq \dim_H(\cdot)$; and Falconer (2003) for more on the Hausdorff dimension and a proof that $\dim_H(X) \leq \dim_B(X)$ in general). Secondly, a set with finite upper box-counting dimension can be embedded into a finite-dimensional Euclidean space using a linear map (a

¹Partially supported by CNPq 302022/2008-2 and 451761/2008-1, CAPES/DGU 267/2008 and FAPESP 2008/53094 and 2008/55516-3, Brazil.

²Partially supported by Ministerio de Ciencia e Innovación grants # MTM2008-0088, # PBH2006-0003-PC, and Junta de Andalucía grants # P07-FQM-02468, # FQM314, Spain.

result which fails for sets that only have finite Hausdorff dimension) – this was originally proved by Mañé in the same paper as the result which forms the main topic of our work here, but his result has been significantly improved in the Hilbert space case by Foias & Olson (1996) [the inverse is Hölder continuous] and Hunt & Kaloshin (1999) [explicit bounds on the Hölder exponent], and in the Banach space case by Robinson (2009) [a Banach space version of Hunt & Kaloshin's result].

We now give a formal definition of the box-counting dimension. Let X be a Banach space and K a compact subset of X . Define $N_X(K, \epsilon)$ as the minimum number of balls in X of radius ϵ needed to cover K . The (upper) box-counting dimension $\dim_B(K)$ of K is defined by:

$$(1.1) \quad \dim_B(K) = \limsup_{\epsilon \rightarrow 0} \frac{\log N_X(K, \epsilon)}{-\log \epsilon}$$

(the lim sup in the definition is necessary; there are sets for which the lim sup is not equal to the lim inf, see for example Mattila, 1995). Essentially this definition extracts the exponent d from the scaling law $N_X(K, \epsilon) \sim \epsilon^{-d}$. More rigorously, $\dim_B(K)$ is the smallest real number such that for any $d > \dim_B(K)$ there exists an $\epsilon_0 > 0$ such that $N_X(K, \epsilon) \leq \epsilon^{-d}$ for all $0 < \epsilon < \epsilon_0$.

The arguments that provide bounds on the dimension of attractors all follow similar lines, which we can formalise in the following lemma.

Lemma 1.1. *Let K be a compact subset of a Banach space X , let $f : X \rightarrow X$ be continuously differentiable in a neighbourhood of K , and let K be negatively invariant for f , i.e. $f(K) \supseteq K$. Suppose that there exist α , $0 < \alpha < 1$ and $M \geq 1$ such that for any $x \in K$,*

$$(1.2) \quad N_X(Df(x)[B_X(0, 1)], \alpha) \leq M.$$

Then

$$(1.3) \quad \dim_B(K) \leq \frac{\log M}{-\log \alpha}.$$

Proof. First, we ensure that (1.2) is sufficient to provide bounds on the number of balls required to cover $f(B_X(x, r))$ when r is small enough. Since f is continuously differentiable and K is compact, for any $\eta > 0$ there exists an $r_0 = r_0(\eta)$ such that for any $0 < r < r_0$ and any $x \in K$,

$$f(B_X(x, r)) \subseteq f(x) + Df(x)[B_X(0, r)] + B_X(0, \eta r),$$

where $A + B$ is used to denote the set $\{a + b : a \in A, b \in B\}$. It follows that

$$(1.4) \quad N_X(f[B_X(x, r)], (\alpha + \eta)r) \leq M$$

for all $r \leq r_0(\eta)$.

Now fix η with $0 < \eta < 1 - \alpha$, and let $r_0 = r_0(\eta)$. Cover K with $N_X(K, r_0)$ balls of radius r_0 . Apply f to every element of this cover. Since $f(K) \supseteq K$, this provides a new cover of K formed by sets all of the form $f(B_X(x, r_0))$, for some $x \in K$. It follows from (1.4) that each of these images can be covered by M balls of radius $(\alpha + \eta)r_0$, ensuring that $N_X(K, (\alpha + \eta)r_0) \leq MN_X(K, r_0)$. Applying this argument k times implies that

$$N_X(K, (\alpha + \eta)^k r_0) \leq M^k N_X(K, r_0).$$

It follows from the definition of $\dim_B(K)$ that

$$\dim_B(K) \leq \frac{\log M}{-\log(\alpha + \eta)},$$

and since $\eta > 0$ we arbitrary we obtain (1.3). $\square$

The key to applying this approach is to be able to prove (1.2), i.e. to find a way of estimating the number of balls of radius α required to cover $Df(x)B(0, 1)$. Our argument is based (as was that of Mallet-Paret and Mañé) on the fact that one can find a sharp estimate for coverings by balls in $(\mathbb{K}^n, \ell^\infty)$, and then use an isomorphism T between $(\mathbb{K}^n, \ell^\infty)$ and an n -dimensional linear subspace U of X to relate coverings with respect to these two different norms. We observe here that only the product $\|T\|\|T^{-1}\|$ occurs in these estimates; this shows that covering results are related to the Banach–Mazur distance between $(U, \|\cdot\|)$ and $(\mathbb{K}^n, \ell^\infty)$, which is bounded by $\log n$ independently of U . Although this follows from the powerful general result that $d_{\text{BM}}(X, Y) \leq \log n$ for any two n -dimensional normed spaces, we give a simple proof of the particular result that we require here.

We use our covering results to prove that a negatively invariant compact set for a nonlinear map with a derivative that is a sum of a strong contraction with a compact map has finite box-counting dimension. Our proof is much simpler than that of Mañé and our bound on the dimension in a Banach space improves on his.

In Section 3 we derive interesting corollaries of the main theorem, and consider some applications. In particular, we show that the global attractors of a wide class of evolution equations are finite dimensional.

2. COVERINGS IN BANACH SPACES

In this section we provide a bound of the form

$$N_X(T[B_X(0, 1)], \alpha) \leq M$$

for linear maps T that are the sum of a compact map and a contraction. When $f : X \rightarrow X$ is such that $Df(x)$ is of this form for all $x \in K$, and the contraction constant is uniformly bounded over K , we show that (1.2) holds uniformly over K .

2.1. The Banach–Mazur distance and coverings of balls in finite-dimensional subspaces. The key result concerns coverings of a ball in a finite-dimensional subspace of X by balls of smaller radius. We use ideas related to the Banach–Mazur distance to provide the estimate we need.

2.1.1. The Banach–Mazur distance. Let X and Y be normed spaces. If there exists $T \in \mathcal{L}(X, Y)$ that is bijective and has $T^{-1} \in \mathcal{L}(Y, X)$, we say that X and Y are isomorphic and that T is an isomorphism between X and Y . The Banach–Mazur distance between two isomorphic normed spaces X and Y , $d_{\text{BM}}(X, Y)$, is defined as

$$\log \left(\inf \left\{ \|T\|_{\mathcal{L}(X, Y)} \|T^{-1}\|_{\mathcal{L}(Y, X)} : T \in \mathcal{L}(X, Y), T^{-1} \in \mathcal{L}(Y, X) \right\} \right).$$

Clearly $d_{\text{BM}}(X, Y) = 0$ if and only if X and Y are isometrically isomorphic; in particular this is the case for any two separable Hilbert spaces of the same cardinality.

It is a consequence of John's Theorem on bounding ellipses of minimal volume (John, 1948) that for any two n -dimensional real Banach spaces U and V , $d_{\text{BM}}(U, V) \leq \log n$, see Bollobás (1990, Theorem 4.15). Here we give a simple proof that if U is any n -dimensional normed space over $\mathbb{K}$ ($= \mathbb{R}$ or $\mathbb{C}$), $d_{\text{BM}}(U, \mathbb{K}_\infty^n) \leq \log n$, where $\mathbb{K}_\infty^n$ denotes $\mathbb{K}^n$ equipped with the ℓ^∞ norm: for $\underline{z} \in \mathbb{K}^n$ with $\underline{z} = (z_1, \dots, z_n)$, $z_j \in \mathbb{K}$, we define

$$\|\underline{z}\|_\infty = \max_{j=1, \dots, n} \|z_j\|_{\mathbb{K}}.$$

In order to show that $d_{\text{BM}}(U, \mathbb{K}_\infty^n) \leq \log n$ we will use an Auerbach basis for X . The proof of the existence of such a basis when X is real is standard (see Bollobás, 1990, Theorem 4.13, for example); we give a proof of the complex case in an appendix.

Lemma 2.1. *Let X be a n -dimensional normed vector space (which may be real or complex). Then, there exists a basis $B = \{x_1, \dots, x_n\}$ for X and a basis $B^* = \{f_1, \dots, f_n\}$ for X^* with $\|x_i\|_X = \|f_i\|_{X^*} = 1$ ($i = 1, \dots, n$) such that $f_i(x_j) = \delta_{ij}$, $i, j = 1, \dots, n$.*

Given this lemma the proof of the following proposition is straightforward.

Proposition 2.2. *Let U be an n -dimensional Banach space over $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$. Then $d_{\text{BM}}(U, \mathbb{K}_\infty^n) \leq \log n$.*

Note that Mañé's paper contains a proof of the bound $d_{\text{BM}}(U, \mathbb{R}_\infty^n) \leq \log(n2^n)$ when U is real.

Proof. Let $\{x_1, \dots, x_n\}$ be an Auerbach basis for U , and $\{f_1, \dots, f_n\}$ the corresponding basis for U^* . Define a map $J : \mathbb{K}_\infty^n \rightarrow U$ by setting

$$J(\underline{z}) = \sum_{j=1}^n z_j x_j.$$

Then

$$\|J(\underline{z})\|_X = \left\| \sum_{j=1}^n z_j x_j \right\|_X \leq \sum_{j=1}^n |z_j| \leq n \|\underline{z}\|_\infty,$$

and so

$$\|J\|_{\mathcal{L}(\mathbb{K}_\infty^n, U)} \leq n.$$

On the other hand, if $x = \sum_{j=1}^n z_j x_j \in U$ with $\|x\|_X \leq 1$ then since $z_j = f_j(x)$,

$$\|J^{-1}(x)\|_\infty = \|\underline{z}\|_\infty = \max_{j=1, \dots, n} |z_j| = \max_{j=1, \dots, n} |f_j(x)| \leq \|x\|_X,$$

which implies that

$$\|J^{-1}\|_{\mathcal{L}(X, X_B^\infty)} \leq 1.$$

□

2.1.2. *The covering lemma.* We use this result to prove our covering lemma:

Lemma 2.3. *If U is an n -dimensional subspace of a real Banach space X , then*

$$(2.1) \quad N_X(B_U(0, r), \rho) \leq (n+1)^n \left(\frac{r}{\rho}\right)^n \quad 0 < \rho \leq r,$$

where the balls in the cover can be taken to have centres in U . The same result holds in a complex Banach space if one replaces the right-hand side of (2.1) with its square.

Proof. Assume first that $\mathbb{K} = \mathbb{R}$. Since U and $\mathbb{R}_\infty^n$ are n -dimensional, $d_{\text{BM}}(U, \mathbb{R}_\infty^n) \leq \log n$: in particular, there exists a linear isomorphism $T : \mathbb{R}_\infty^n \rightarrow U$ such that $\|T\|\|T\|^{-1} \leq n$. Since

$$B_U(0, r) = TT^{-1}(B_U(0, r)) \subseteq T(B_{\mathbb{R}_\infty^n}(0, \|T^{-1}\|r)),$$

and $B_{\mathbb{R}_\infty^n}(0, \|T^{-1}\|r)$ can be covered by

$$\left(1 + \frac{\|T^{-1}\|r}{\rho/\|T\|}\right)^n = \left(1 + \|T\|\|T^{-1}\|\frac{r}{\rho}\right)^n \leq \left(1 + n\frac{r}{\rho}\right)^n \leq (n+1)^n \left(\frac{r}{\rho}\right)^n$$

balls in $\mathbb{R}_\infty^n$ of radius $\rho/\|T\|$, it follows that $B_U(0, r)$ can be covered by the same number of U -balls of radius ρ . If X is complex one requires $(1 + (a/b))^{2n}$ b -balls in $\mathbb{C}_\infty^n$ to cover a ball of radius a . $\square$

2.2. Coverings of $T[B_X(0, 1)]$ via finite-dimensional approximations. We now have good estimates for the coverings of balls in finite-dimensional linear subspaces, but we want to cover the images of balls under linear maps. In order to do this we show that, given a linear map T that is the sum of a compact map and a contraction, $T[B_X(0, 1)]$ can be well-approximated by $T[B_Z(0, 1)]$, where Z is a finite-dimensional subspace of X .

We denote by $\mathcal{L}(X)$ the space of bounded linear transformations from X into itself, by $\mathcal{K}(X)$ the closed subspace of $\mathcal{L}(X)$ consisting of all compact linear transformations from X into itself, and define

$$(2.2) \quad \mathcal{L}_\lambda(X) = \{T \in \mathcal{L}(X) : T = L + C, \text{ with } C \in \mathcal{K}(X) \text{ and } \|L\|_{\mathcal{L}(X)} < \lambda\}.$$

By $\text{dist}(A, B)$ we denote the Hausdorff semi-distance between A and B ,

$$\text{dist}(A, B) = \sup_{a \in A} \left(\inf_{b \in B} \|a - b\|_X \right).$$

Lemma 2.4. *Let X be a Banach space and $T \in \mathcal{L}_{\lambda/2}(X)$. Then there exists a finite-dimensional subspace Z of X such that*

$$(2.3) \quad \text{dist}(T[B_X(0, 1)], T[B_Z(0, 1)]) < \lambda.$$

We denote by $\nu_\lambda(T)$ the minimum $n \in \mathbb{N}$ such that (2.3) holds for some n -dimensional subspace of X .

Proof. Write $T = L + C$, where $C \in \mathcal{K}(X)$ and $L \in \mathcal{L}(X)$ with $\|L\|_{\mathcal{L}(X)} < \lambda/2$. We show first that for any $\epsilon > 0$ there is a finite-dimensional subspace Z such that

$$(2.4) \quad \text{dist}(C[B_X(0, 1)], C[B_Z(0, 1)]) < \epsilon.$$

Suppose that this is not the case. Choose some $x_1 \in X$ with $\|x_1\|_X = 1$, and let $Z_1 = \text{span}\{x_1\}$. Then

$$\text{dist}(C[B_X(0, 1)], C[B_{Z_1}(0, 1)]) \geq \epsilon,$$

and so there exists an $x_2 \in X$ with $\|x_2\|_X = 1$ such that

$$\|Cx_2 - Cx_1\|_X \geq \epsilon.$$

With $Z_2 = \text{span}\{x_1, x_2\}$, one can find an x_3 with $\|x_3\|_X = 1$ such that

$$\|Cx_3 - Cx_1\|_X \geq \epsilon \quad \text{and} \quad \|Cx_3 - Cx_2\|_X \geq \epsilon.$$

Continuing inductively one can construct in this way a sequence $\{x_j\}$ with $\|x_j\| = 1$ such that

$$\|Cx_i - Cx_j\|_X \geq \epsilon \quad i \neq j,$$

contradicting the compactness of C .

Now let $\tilde{\lambda} < \lambda$ be such that $2\|L\|_{\mathcal{L}(X)} < \tilde{\lambda} < \lambda$, and choose Z using the above argument so that

$$\text{dist}(C[B_X(0, 1)], C[B_Z(0, 1)]) < \lambda - \tilde{\lambda}.$$

If $x \in B_X(0, 1)$ and $z \in B_Z(0, 1)$, then

$$\|Tx - Tz\|_X \leq \|L(x - z)\|_X + \|Cx - Cz\|_X \leq \tilde{\lambda} + \|Cx - Cz\|_X$$

Hence,

$$\begin{aligned} \text{dist}(T[B_X(0, 1)], T[B_Z(0, 1)]) &\leq \tilde{\lambda} + \text{dist}(C[B_X(0, 1)], C[B_Z(0, 1)]) \\ &< \lambda. \end{aligned}$$

This completes the proof. □

2.3. Uniform estimates for $x \in K$.

Theorem 2.5 (after Mañé, 1981). *Let X be a Banach space, $U \subset X$ an open set, and $f : U \rightarrow X$ a continuously differentiable map. Suppose that K is a compact set and assume that for some λ with $0 < \lambda < \frac{1}{2}$,*

$$Df(x) \in \mathcal{L}_{\lambda/2}(X) \quad \text{for all } x \in K.$$

Then $n = \sup_{x \in K} \nu_\lambda(Df(x))$ and $D = \sup_{x \in K} \|Df(x)\|$ are finite, and

$$(2.5) \quad N(Df(x)[B_X(0, 1)], 2\lambda) \leq \left[(n+1) \frac{D}{\lambda} \right]^{\alpha n} \quad \text{for all } x \in K,$$

where $\alpha = 1$ if X is real and $\alpha = 2$ if X is complex. It follows that

$$(2.6) \quad \dim_B(K) \leq \alpha n \left\{ \frac{\log((n+1)D/\lambda)}{-\log(2\lambda)} \right\},$$

Proof. First we show that $n = \sup_{x \in K} \nu_\lambda(Df(x))$ is finite. For each $x \in K$, there exists a finite-dimensional linear subspace Z_x such that

$$\text{dist}(Df(x)[B_X(0, 1)], Df(x)[B_{Z_x}(0, 1)]) < \lambda.$$

Since $Df(\cdot)$ is continuous, it follows that there exists a $\delta_x > 0$ such that

$$\text{dist}(Df(y)[B_X(0, 1)], Df(y)[B_{Z_x}(0, 1)]) < \lambda$$

for all $y \in B_X(x, \delta_x)$, i.e. $\nu_\lambda(y) \leq \nu_\lambda(x)$ for all such y . The open cover of K formed by the union of $B_X(x, \delta_x)$ over x has a finite subcover, from whence it follows that $n < \infty$.

Now, since $n = \sup_{x \in K} \nu_\lambda(Df(x)) < \infty$, for each $x \in K$ there is a subspace Z_x of X with $\dim(Z_x) \leq n$ such that

$$\text{dist}(Df(x)[B_X(0, 1)], Df(x)[B_{Z_x}(0, 1)]) < \lambda.$$

For ease of notation we now drop the x subscript on Z_x , and write $T = Df(x)$.

Noting that $T(Z)$ is also an n -dimensional subspace of X , one can use Lemma 2.3 to cover the ball $B_{T(Z)}(0, \|T\|)$ with balls $B_X(y_i, \lambda)$, $1 \leq i \leq k$, such that $y_i \in B_X(0, \|T\|)$ for each i and

$$k \leq \left[(n+1) \frac{\|T\|}{\lambda} \right]^{\alpha n}.$$

Thus

$$(2.7) \quad T[B_Z(0, 1)] \subseteq B_{T(Z)}(0, \|T\|) = B_X(0, \|T\|) \cap T(Z) \subseteq \bigcup_{i=1}^k B_X(y_i, \lambda).$$

We complete the proof by showing that

$$\bigcup_{i=1}^k B_X(y_i, 2\lambda) \supseteq T[B_X(0, 1)].$$

Indeed, if $x \in B_X(0, 1)$ then it follows from (2.3) that there is a $y \in T[B_Z(0, 1)]$ such that $\|Tx - y\|_X < \lambda$. Since $y \in T[B_Z(0, 1)]$, it follows from (2.7) that $\|y - y_i\|_X \leq \lambda$ for some $i \in \{1, \dots, k\}$, and so

$$\|Tx - y_i\|_X \leq \|Tx - y\|_X + \|y - y_i\|_X < 2\lambda,$$

i.e. $x \in B_X(y_i, 2\lambda)$.

The result now follows as stated since n is uniform over $x \in K$. $\square$

Some immediate improvement of the above is possible if we work in a Hilbert space. A result due to Cheyphzhov & Vishik (2002) (Lemma III.2.1) shows that in $\mathbb{R}^n$ we need no more than $4^n(r/\rho)^n$ balls of radius ρ to cover a ball of radius $r > \rho$. This implies that one can replace the factor $(n+1)^n$ in (2.1) by 7^n , and obtain a corresponding improvement in the bound of Theorem 2.5. We do not pursue this direction further here, since the use of ellipses in place of balls (an approach initiated by Douady & Oesterlé, 1980, and developed further by Constantin & Foias, 1985) leads to significant improvements on the possible dimension estimates and is now the standard approach.

3. COROLLARIES & APPLICATIONS

3.1. When $Df \in \mathcal{L}_1(X)$. The following corollary can be found in Hale, Maghalaes, & Oliva (2002):

Corollary 3.1. *Suppose that X is a Banach space, $U \subset X$ an open set, and $f : U \rightarrow X$ a continuously differentiable map. Suppose that $K \subset U$ is a compact set such that $f(K) \supseteq K$, and that $Df(x) \in \mathcal{L}_1(X)$ for all $x \in K$. Then $\dim_B(K) < \infty$.*

Proof. It follows from an argument similar to that used in the proof of Theorem 2.5 to show that $n < \infty$ that in fact there exists an $\alpha < 1$ such that $Df(x) \in \mathcal{L}_\alpha(X)$ for all $x \in K$. Note that

$$D[f^p] = Df(f^{p-1}(x)) \circ \cdots \circ Df(x),$$

and that if $C_i \in \mathcal{K}(X)$ and $L_i \in \mathcal{L}(X)$, $i = 1, 2$, then

$$(C_1 + L_1) \circ (C_2 + L_2) = \underbrace{[C_1 \circ C_2 + C_1 \circ L_2 + L_1 \circ C_2]}_{\in \mathcal{K}(X)} + L_1 \circ L_2,$$

it follows that if $Df(x) \in \mathcal{L}_\alpha(X)$ with $\alpha < 1$ then $[D(f^p)](x) \in \mathcal{L}_{\alpha^p}(X)$. It follows that for p large enough, $D(f^p)(x) \in \mathcal{L}_\lambda$ for some $\lambda < 1/4$, for every $x \in K$. One can now apply Theorem 2.5 to f^p in place of f (noting that $f^p(K) \supseteq K$) to deduce that $d_f(K) < \infty$. $\square$

3.2. When $D_x T$ has finite rank.

Corollary 3.2. *Let X be a Banach space and assume that $T \in \mathcal{C}^1(X)$, K is a compact set such that $T(K) = K$, and $D_x T$ has finite rank $\nu(x)$ with $\sup_{x \in K} \nu(x) := \nu < \infty$. Then,*

$$\dim_B(K) \leq \nu.$$

Proof. Clearly, for each $\lambda > 0$ and $x \in K$, $D_x T \in \mathcal{L}_{\frac{\lambda}{2}}(X)$ for all $\lambda > 0$. Consequently, for each $0 < \lambda < \frac{1}{2}$,

$$\dim_B(K) \leq \nu \frac{\log((\nu + 1)\frac{D}{\lambda})}{\log(1/2\lambda)}.$$

Taking the limit as $\lambda \rightarrow 0$ we have that $\dim_B(K) \leq \nu$. $\square$

3.3. An ordinary differential equation.

Corollary 3.3. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuously differentiable function. Assume that the semigroup $\{S(t) : t \geq 0\}$ in $\mathbb{R}^n$ associated to the ordinary differential equation*

$$\dot{x} = f(x) \quad x(0) = x_0 \in \mathbb{R}^n.$$

has a global attractor $\mathcal{A}$. If $\text{rank}(D_x f) \leq k \leq n$ for all $x \in \mathcal{A}$, then $\dim_B(\mathcal{A}) \leq k$.

In particular, if $f : \mathbb{R}^k \rightarrow \mathbb{R}^k$, $\beta > 0$ and there is a constant $M > 0$ such that $f(x) \cdot x < 0$ for $\|x\|_{\mathbb{R}^k} \geq M$, then the semigroup $\{S(t) : t \geq 0\}$ associated to

$$\begin{aligned} \frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 & I \\ 0 & -\beta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ f(x) \end{pmatrix} \\ \begin{pmatrix} x \\ y \end{pmatrix} (0) &= \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \end{aligned}$$

has a global attractor $\mathcal{A}$ in $\mathbb{R}^k \times \mathbb{R}^k$ with $\dim_B(\mathcal{A}) \leq k$.

3.4. Semilinear equations in Banach spaces.

Corollary 3.4. *Let $A : D(A) \subset X \rightarrow X$ be a sectorial operator with $\operatorname{Re} \sigma(A) > 0$. If $f : X^\alpha \rightarrow X$ is continuously differentiable and Lipschitz continuous in bounded subsets of X^α and the semigroup $\{S(t) : t \geq 0\}$ in X^α associated to the abstract parabolic problem*

$$\dot{x} + Ax = f(x) \quad \text{with} \quad x(0) = x_0 \in X^\alpha$$

has a global attractor $\mathcal{A}$ and either e^{-At} is compact for each $t > 0$ or $f_x \in \mathcal{K}(X^\alpha, X)$ is compact for each $x \in \mathcal{A}$, then $\dim_B(\mathcal{A}) < \infty$.

Proof. For $x \in \mathcal{A}$, let

$$S(t)x = e^{-At}x + \int_0^t e^{-A(t-s)}f(S(s)x)ds$$

hence, the derivative $S_x(t) \in \mathcal{L}(X^\alpha)$ with respect to x of $S(t)$ at x satisfies

$$S_x(t) = e^{-At} + \int_0^t e^{-A(t-s)}f'(S(s)x)S_x(s)ds.$$

Hence, for t suitably large, the hypothesis of Corollary 3.1 are satisfied and the result follows. $\square$

Now we show how a rough estimate on the dimension of the attractor can be obtained. First note that, if $A : D(A) \subset X \rightarrow X$ is a sectorial operator with compact resolvent, X^β , $\beta \geq 0$ denotes the fractional power spaces associated to A , $\alpha > 0$, there is a sequence of finite rank projections $\{P_n\}_{n \in \mathbb{N}}$ and sequences of positive real numbers $\{\lambda_n\}_{n \in \mathbb{N}}$ and $\{M_n\}_{n \in \mathbb{N}}$ such that

$$(3.1) \quad \|e^{-At}(I - P_n)\|_{\mathcal{L}(X^\gamma, X^\beta)} \leq M_n t^{-(\beta-\gamma)} e^{-\lambda_n t}, \quad t \geq 0, \quad 0 \leq \gamma \leq \beta \leq \alpha.$$

We say that A is an admissible sectorial operator if it is sectorial and there is a sequence $\{\lambda_n\}_{n \in \mathbb{N}}$ and $M > 0$ such that (3.1) with $M_n = M$ for all $n \in \mathbb{N}$. It is not difficult to see that, if A is an admissible sectorial operator, then A has compact resolvent.

Also

$$\|S_x(t)\|_{\mathcal{L}(X^\alpha)} \leq M + MN \int_0^t (t-s)^{-\alpha} \|S_x(s)\|_{\mathcal{L}(X^\alpha)} ds,$$

where

$$N = \sup \{ \|f'(x)\|_{\mathcal{L}(X^\alpha, X)} : x \in \mathcal{A} \}$$

It follows from the singular Gronwall's inequality (see Lemma 7.1.2 in Henry 1980) that

$$\|S_x(t)\|_{\mathcal{L}(X^\alpha)} \leq \frac{\bar{M}}{1-\alpha} e^{(MN\Gamma(1-\alpha))^{1/(1-\alpha)} t}.$$

Now, if $Q_n = (I - P_n)$,

$$\|Q_n T_x(t)\|_{\mathcal{L}(X^\alpha)} \leq M e^{-\lambda_n t} + MN \int_0^t (t-s)^{-\alpha} e^{-\lambda_n(t-s)} \|S_x(s)\|_{\mathcal{L}(X^\alpha)} ds$$

and

$$\begin{aligned}
& \|Q_n T_x(t)\|_{\mathcal{L}(X^\alpha)} \\
& \leq M e^{-\lambda_n t} + \frac{\bar{M} M N}{1 - \alpha} e^{(M N \Gamma(1-\alpha))^{1/(1-\alpha)}} \int_0^t (t-s)^{-\alpha} e^{-(\lambda_n + (M N \Gamma(1-\alpha))^{1/(1-\alpha)})(t-s)} ds \\
& \leq M e^{-\lambda_n t} + \frac{\bar{M} M N}{1 - \alpha} e^{(M N \Gamma(1-\alpha))^{1/(1-\alpha)} t} \int_0^t u^{-\alpha} e^{-(\lambda_n + (M N \Gamma(1-\alpha))^{1/(1-\alpha)})u} du \\
& \leq M e^{-\lambda_n t} + \frac{\bar{M} M N}{1 - \alpha} \frac{e^{(M N \Gamma(1-\alpha))^{1/(1-\alpha)} t} \Gamma(1-\alpha)}{\lambda_n + (M N \Gamma(1-\alpha))^{1/(1-\alpha)}} = \Lambda_n(t), \quad t \geq 0.
\end{aligned}$$

From the admissibility of A and (3.1), $\|Q_n\|_{\mathcal{L}(X^\alpha)} \leq M$, for all $n \in \mathbb{N}$. Hence,

$$\|Q_n T_x(t) Q_n\|_{\mathcal{L}(X^\alpha)} \leq M \Lambda(t), \quad t \geq 0.$$

Choose $t = 1$ and $n_0 \in \mathbb{N}$ such that $\Lambda(1) < \lambda < \frac{1}{4}$. If $F = S(1)$, $L = Q_{n_0} S(1)$ and $C = P_{n_0} S(1)$, then $\mathcal{A}$ is invariant under F . Furthermore, $F_x = L_x + C_x$ with $L_x = Q_{n_0} S_x(1)$ and $C_x = P_{n_0} S_x(1)$ and if $Z_x = R(C_x)$ and W_x is a subspace of X such that $C_x : W_x \rightarrow Z_x$ is an isomorphism, $F_x \in \mathcal{L}_{\lambda/2}(X)$ for all $x \in \mathcal{A}$ and for some $\lambda < \frac{1}{2}$. In addition,

$$\nu = \sup_{x \in \mathcal{A}} \dim(Z_x) \leq \dim(R(P)).$$

This proves that all the assumptions of Theorem 2.5 are satisfied and consequently

$$(3.2) \quad \dim_B(\mathcal{A}) \leq \nu \frac{\log((\nu + 1) \frac{D}{\lambda})}{\log(1/2\lambda)} < \infty.$$

4. CONCLUSION

As specific examples of the result in Section 3 (a) we mention the attractor for the damped wave equation with critical exponent (see Arrieta et al (1992)) and as an application of the results in Section 3 (d) we mention the attractors of dissipative parabolic equations in $L^p(\Omega)$, $W^{1,p}(\Omega)$ as in Arrieta et al (2000) or the Navier-Stokes Equation in space dimension 2 as in Temam (1988).

As an example of a problem which does not define a semigroup in a Hilbert space we mention that of Arrieta et al (submitted for publication), for Section 3 (d). For Section 3 (a), the examples are the attractors for functional differential equations for which the natural phase space is not a Hilbert space (see Hale et al., 2002, for example).

APPENDIX: EXISTENCE OF AN AUERBACH BASIS IN A COMPLEX BANACH SPACE

In this appendix we give a proof of the existence of an Auerbach basis in a complex n -dimensional Banach space. The result is standard in a real Banach space, and we use the real version in our proof.

Lemma 4.1. *Let X be a n -dimensional complex normed vector space. Then, there exists a basis $B = \{x_1, \dots, x_n\}$ for X and a basis $B^* = \{f_1, \dots, f_n\}$ for X^* with $\|x_i\|_X = \|f_i\|_{X^*} = 1$ ($i = 1, \dots, n$) such that $f_i(x_j) = \delta_{ij}$, $i, j = 1, \dots, n$.*

Proof. Given a basis $B_0 = \{y_1, \dots, y_n\}$ of X we consider the real space X_2 given as the linear span of over $\mathbb{R}$ of

$$B_2 = \{y_1, \dots, y_n, iy_1, \dots, iy_n\},$$

equipped with the norm $\|z\|_{X_2} = \|z\|_X$. We now apply the real version of the result to X_2 , to produce a basis $\{x_1, \dots, x_{2n}\}$ for X_2 and $\{\varphi_1, \dots, \varphi_{2n}\}$ for X_2^* such that $\varphi_j(x_k) = \delta_{jk}$ and $\|\varphi_j\| = \|x_j\| = 1$.

Since $x_j \in X$ and the $\{x_j\}$ must span X over $\mathbb{C}$, we can reorder and relabel the $\{x_j\}$ (and the corresponding $\{\varphi_j\}$) so that $\{x_1, \dots, x_n\}$ span X over $\mathbb{C}$. It follows that

$$\{x_1, \dots, x_n, ix_1, \dots, ix_n\}$$

span X over $\mathbb{R}$. Furthermore, each ix_j must be a linear combination of $\{x_{n+1}, \dots, x_{2n}\}$, since it follows from the fact that $\{x_1, \dots, x_n\}$ form a basis for X over $\mathbb{C}$ that $\{ix_1, x_2, x_n\}$ are linearly independent (over $\mathbb{C}$, so certainly over $\mathbb{R}$).

For $k = 1, \dots, n$ we define an element $f_k \in X^*$ via

$$f_k(z) = \varphi_k(z) - i\varphi_k(iz) \quad z \in X,$$

where in order to interpret $\varphi_k(z)$ we consider z as an element of X_2 (expand in terms of the basis $\{x_1, \dots, x_n\}$ and split the real and imaginary parts of the coefficients). We now show that the $\{f_k\}$ have the properties we require.

First, note that for $1 \leq k, j \leq n$,

$$f_k(x_j) = \varphi_k(x_j) - i\varphi_k(ix_j) = \delta_{kj},$$

since ix_j is a linear combination of $\{x_{n+1}, \dots, x_{2n}\}$, and thus $\varphi_k(ix_j) = 0$. All that remains is to show that $\|x_k^*\| = 1$. To do this we follow the standard argument (e.g. Yosida, 1980), writing $f_k(z) = re^{-i\theta}$. Then

$$|f_k(z)| = e^{i\theta} f_k(z) = f_k(e^{i\theta} z),$$

so that $f_k(e^{i\theta} z)$ is real and positive. It follows that

$$|f_k(z)| = f_k(e^{i\theta} z) = \varphi_k(e^{i\theta} z) \leq \|e^{i\theta} z\|_{X_2} \leq \|e^{i\theta} z\|_X = \|z\|_X,$$

and the lemma is proved. $\square$

Acknowledgement: Part of this work has been carried out while ANC and JAL were visiting the Warwick Mathematics Institute, University of Warwick, UK. They both wish to acknowledge the great hospitality of the people in this institution. The paper was completed while all three authors were visiting the de Giorgi Centre in Pisa; we are very grateful for the opportunity this visit gave us to continue our collaboration. ANC was partially supported by grants CNPq 302022/2008-2 and 451761/2008-1, CAPES/DGU 267/2008 and FAPESP 2008/53094 and 2008/55516-3, Brazil; JAL by Ministerio de Ciencia e Innovación grant MTM2008-0088, and Junta de Andalucía grants P07-FQM-02468 and FQM314; JCR wishes to acknowledge the support of the EPSRC, grant EP/G007470/1.

REFERENCES

- [1] J. M. Arrieta, A.N. Carvalho and A. Rodriguez-Bernal, Attractors for Parabolic Problems with Nonlinear Boundary Condition. Uniform Bounds, *Communications in Partial Differential Equations* **25** (1-2), 1-37 (2000).
- [2] J. M. Arrieta, A.N. Carvalho and J. K. Hale, A Damped Hyperbolic Equation with Critical Exponent. *Communications in Partial Differential Equations*, **17** (5-6) 841-866 (1992).
- [3] J. M. Arrieta, A.N. Carvalho and G. Lozada-Cruz, Dynamics in dumbbell domains II. The Limiting Problem, submitted for publication.
- [4] B. Bollobás, *Linear Analysis: Introductory Course*, Cambridge Mathematical Textbooks, Cambridge University Press, Cambridge UK (1990)
- [5] V.V. Chepyzhov and M. I. Vishik, *Attractors for equations of mathematical physics*. American Mathematical Society Colloquium Publications, **49**. American Mathematical Society, Providence, RI, 2002.
- [6] P. Constantin and C. Foias, Global Lyapunov exponents, Kaplan-Yorke formulas and the dimension of the attractors for 2D Navier-Stokes equations. *Comm. Pure Appl. Math.* **38** (1985), 1-27.
- [7] A. Douady and J. Oesterlé, Dimension de Hausdorff des attracteurs. *C. R. Acad. Sci. Paris Sér. A* **290** (1980), 1135-1138.
- [8] K. Falconer, *Fractal geometry. Mathematical foundations and applications*. Second edition. John Wiley & Sons, Inc., Hoboken, NJ, 2003.
- [9] C. Foias and E. J. Olson, *Finite fractal dimension and Hölder-Lipschitz parametrization*. *Indiana Univ. Math. J.* **45** (1996), 603-616.
- [10] J. K. Hale, L. T. Magalhães and W. M. Oliva, *Dynamics in infinite dimensions. With an appendix by Krzysztof P. Rybakowski*. Second edition. Applied Mathematical Sciences, **47**. Springer-Verlag, New York, 2002.
- [11] B. R. Hunt and V. Y. Kaloshin, *Regularity of embeddings of infinite-dimensional fractal sets into finite-dimensional spaces*. *Nonlinearity* **12** (1999), 1263-1275.
- [12] D. Henry, *Geometric theory of semilinear parabolic equations*, Lecture Notes in Mathematics **840**, Springer-Verlag (1981).
- [13] W. Hurewicz and H. Wallman, *Dimension Theory*. Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941.
- [14] F. John, Extremum problems with inequalities as subsidiary conditions. Studies and Essays Presented to R. Courant on his 60th Birthday, January 8, 1948, 187-204. Interscience Publishers, Inc., New York, N. Y., 1948.
- [15] J. Mallet-Paret, *Negatively invariant sets of compact maps and an extension of a theorem of Cartwright*, *J. Differential Equations*, **22**, 331-348 (1976).
- [16] R. Mañé, *On the dimension of the compact invariant sets of certain nonlinear maps*. Springer Lecture Notes in Math., vol. 898, Springer, New York, 230-242, 1981.
- [17] P. Mattila, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*. Cambridge Studies in Advanced Mathematics, **44**. Cambridge University Press, Cambridge, 1995.
- [18] J. C. Robinson, *Linear embeddings of finite-dimensional subsets of Banach spaces into Euclidean spaces*. *Nonlinearity* **22** (2009), 711-728.
- [19] R. Temam, *Infinite-dimensional dynamical systems in mechanics and physics*. Applied Mathematical Sciences, **68**. Springer-Verlag, New York, 1988 (Second edition, 1997).

(A. N. Carvalho) INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO-CAMPUS DE SÃO CARLOS, CAIXA POSTAL 668, 13560-970 SÃO CARLOS SP, BRAZIL

(J. A. Langa) DEPARTAMENTO DE ECUACIONES DIFERENCIALES Y ANÁLISIS NUMÉRICO, UNIVERSIDAD DE SEVILLA, APDO. DE CORREOS 1160, 41080-SEVILLA SPAIN

(J. C. Robinson) MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL. U.K.

NOTAS DO ICMC

SÉRIE MATEMÁTICA

- 305/09 CARABALLO, T.; CARVALHO, A.N.; LANGA, J.A.; RIVERO, F. – Existence of pullback attractors for pullback asymptotically compact processes.
- 304/09 CARABALLO, T.; LANGA, J.A.; RIVERO, F.; CARVALHO, A.N. – A gradient-like non-autonomous evolution process.
- 303/09 CAMPOS, J.E.P.P. – Distinguishing link-homotopy classes by algebraic methods.
- 302/08 RODRIGUES, H.M.; SOLÀ-MORALES, J. – A note on the relationship between spectral radius and norms of bounded linear operators.
- 301/08 ARRIETA, J.M.; CARVALHO, A.N.; LOZADA-CRUZ, G. – Dynamics in dumbbell domains III. Continuity of attractors.
- 300/08 ARRIETA, J.M.; CARVALHO, A.N.; LOZADA-CRUZ, G. – Dynamics in dumbbell domains II. The limiting problem.
- 299/08 DEMUNER, D.P.; FEDERSON, M.; GUTIERREZ, C. – The Poincaré-Bendixson theorem on the Klein bottle for continuous vector fields.
- 298/08 BONOTTO, E.M.; GIMENES, L.P.; FEDERSON, M.; – Oscillation for a second-order neutral differential equation with impulses.
- 297/08 CALLEJAS-BEDREGAL, R.; JORGE PÉREZ, V.H. – Rees's mixed multiplicity theorem for modules.
- 296/08 BONOTTO, E.M. – LaSalle's theorems in impulsive semidynamical systems.